



## NEBRASKA'S WATER MANAGEMENT RESOURCE

Providing the sound science and support for managing  
Nebraska's most precious resource.

# **Classification of Irrigated and Non-Irrigated Land Using Remote Sensing Techniques: *A Case Study in Nebraska***

**Mahesh Pun, EIT**

**Ruopu Li, PhD**

Integrated Water Management Analyst

Nebraska Department of Natural Resources


2015 NGWA Conference on the Upper Great Plains

Cayenne, Wyoming

September 22, 2015

# Irrigated Agriculture in Nebraska



- Agriculture plays a pivotal role in Nebraska economy
- Nebraska ranks 1st in irrigated acres



Source: USDA (2013)

# Irrigated/Non-irrigated Farmland


- Irrigated Farmland
  - Irrigation meets the crop needs when lack of rain during the growing season
- Non-Irrigated Farmland
  - Only rain-fed crops
  - Susceptible to drought



Source: <http://www.fao.org/docrep/006/y4683e/y4683e07.htm>

# Impacts of Irrigation

- Managing Impacts of Irrigation




# Project Goals

- To develop a scientifically defensible and cost-effective technique for classifying irrigated and non-irrigated farmland using remote sensing techniques

# Methods

- Data
  - Landsat Remote Sensing Imagery
- Stage 1
  - Development of irrigated land area based on pixel-based classification
- Stage 2
  - Development of irrigated land area using object-oriented classification
- Stage 3
  - Automate the entire process

# Study Area





# Methods—Pixel-based Classification

- Normalized Difference Vegetation Index (NDVI)
  - Popular vegetation and irrigation monitoring tool
- Greenness Index (GI)
  - Sensitive to soil moisture stress than NDVI
- Evaporative fraction (ETRF)
  - Indicating water stress; more responsive than NDVI
  - Surface Energy Balance System (SEBS)

$$ETRF = \frac{\lambda E}{Rn - G}$$



# Methods—Pixel-based Classification


- Two new index
  - Enhance the spectral contrast


$$NGI = NDVI * GI$$

$$EGI = \frac{EFRF}{GI}$$

- Calibration with Ground-truth data
- Verification with the NASS data


$$MAPE = \frac{1}{N} \sum \left( \frac{|NASS - NEG|}{NASS} \right)$$






# Methods—Object-oriented Classification

- Recently developed classification method working on image objects rather than pixels
- What is object?
  - A cluster of adjacent pixels with similar spectral values
  - Can be linked with real ground objects
  - Allows a rich collection of descriptors
    - Such as texture, color, shape, topology
- eCognition Software
  - Rich set of tools working with objects





# Methods—Object-oriented Classification

- Image segmentation
  - Grouping pixels into objects!
- Image classification
  - Classification of objects
  - A combination of methods
- Classification refinement
- Vectorization
  - Output as vector data






# Results—Pixel-based Classification



# Results—Pixel-based Classification

- Verification with USDA NASS irrigated acres




| COUNTY   | NASS   | NEG      | % Error | NASS   | NEG      | % Error |
|----------|--------|----------|---------|--------|----------|---------|
| YEAR     | 2014   |          |         | 2012   |          |         |
| Adams    | 222400 | 206452.8 | -7.2    | 225700 | 199703.3 | -11.5   |
| Clay     | 214000 | 219221.8 | 2.4     | 211900 | 223073.7 | 5.3     |
| Fillmore | 232400 | 246762.3 | 6.2     | 226300 | 240420.2 | 6.2     |
| Franklin | 93500  | 94689.35 | 1.3     | 101600 | 110192.9 | 8.5     |
| Harlan   | 49000  | 52035.87 | 6.2     | 92700  | 92546.36 | -0.2    |
| Kearney  | 192200 | 191280.2 | -0.5    | 215700 | 228001.1 | 5.7     |
| Nuckolls | 61100  | 74725.22 | 22.3    | 67300  | 86034.42 | 27.8    |
| Thayer   | 162300 | 180392.9 | 11.1    | 153600 | 180392.9 | 17.4    |
| Webster  | 51200  | 52499.34 | 2.5     | 62300  | 69067.2  | 9.8     |
| Furnas   | 39400  | 43389.83 | 10.1    | 54100  | 51665.36 | -4.5    |
| Phelps   | *      | 232239.9 | -       | 246200 | 245411.2 | -0.3    |
| Gosper   | *      | 79196.6  | -       | 86700  | 83523.59 | -3.7    |



# Results—Object-oriented Classification

Nuckolls County

| Classification | NASS  | Difference |
|----------------|-------|------------|
| 63624          | 61100 | 4%         |







## Discussion

- **ET** is an important component for differentiating spectral signature of irrigated/non-irrigated fields even during the wet year (2014)
- The pixel-based classification can produce results **comparable** with NASS county data
- The object-oriented approach can help **refine** the results and automate the work flow
- The method and results can help **improve** water resource decision making

# Discussion

- Limitation
  - Assuming the NASS county-level irrigated acres are 'accurate'
  - The spatial distribution of ground truth was limited by the accessibility of field crews and funding
  - Cloud and scan-line gaps are interpolated



## Future Work

- Comparing pixel-based and object-oriented classification results for each county
- Calibrating with better ground-truth data
- Building an automated work flow
  - Combining two methods into one
- Auxiliary data can be used for refinement

# Summary

- This study shows the advantages of remote sensing techniques for estimating irrigated and non-irrigated fields
- Better decisions on water resource management can be made with this method
- More work will be done to integrate and automate the entire work flow

# Any Questions?



Department of Natural Resources

NEBRASKA'S WATER MANAGEMENT RESOURCE

Providing the sound science and support for managing  
Nebraska's most precious resource.

**THANK YOU**

**Mahesh Pun, EIT**

Integrated Water Management Analyst

[Mahesh.pun@nebraska.gov](mailto:Mahesh.pun@nebraska.gov)

402-471-3959

Nebraska Department of Natural Resources

402-471-2363

[dnr.nebraska.gov](http://dnr.nebraska.gov)

